

Axionic Inflation from Large Volume Flux Compactifications

R. Holman* and Jimmy A. Hutasoit†

Dept. of Physics, Carnegie Mellon University, Pittsburgh, PA 15213, USA.

(Dated: February 1, 2008)

We find a general model of *single-field* inflation within the context of type IIB string theory compactified on large volume Calabi-Yau orientifolds with $h^{2,1} > h^{1,1} = 2$. The inflaton is the axion part of the complexified Kähler moduli and the resulting scalar power spectrum is red, and can easily be made compatible with WMAP3 bounds on n_s . This model overcomes the η -problem using gaugino condensates on wrapped D7-brane while keeping the tuning of the parameters minimal.

PACS numbers: 11.25.Mj, 11.25.Wx, 98.80.Cq, 98.80.-k

The recent advances in using both NS-NS and R-R fluxes to stabilize the moduli of Type IIB Calabi-Yau (CY) compactifications [1, 2, 3] have sparked intense interest in the possibility of generating the scalar potentials required for inflation (for previous attempts at string inflation, see [4]). This has already resulted in some interesting models, such as racetrack inflation [5], which use a combination of flux generated potentials together with non-perturbative potentials to construct a multi-field inflationary model.

Building on the work of Ref.[2] (KKLT) and Balasubramanian *et al.* [6] (BBCQ), we show here that a *single-field* inflationary model can be obtained from Type IIB compactifications on CY orientifolds in the large volume limit. The inflaton field is one of the axion fields arising as the real part of a complexified Kähler modulus of the compactification. This model solves the η -problem of supergravity inflation models in a novel way: the smallness of η comes from the size of a gauge group $SU(N)$ on wrapped D7-brane that gives rise to gaugino condensation.

The main idea is the following. BBCQ showed that in the large volume limit and neglecting exponentially-suppressed contributions to the potential, the potential has a non-supersymmetric minimum so that one can stabilize the volume and some of the Kähler moduli. They also argued that since this minimum is tachyon-free, one should be able to fix all Kähler and complex structure moduli.

We proceed from where BBCQ left off and go back to the exponentially-suppressed parts of the potential. This potential will stabilize the remaining Kähler moduli and we will find that for Calabi-Yau orientifolds with $h^{1,1} = 2$ (e.g.: $\mathbb{P}^4_{[1,1,1,6,9]}$ [7]), this potential will give rise to inflation.

This model can be thought of as a string version of natural inflation models [8] that used Pseudo-Nambu-Goldstone bosons as inflatons.

Let us first review the relevant aspects of moduli stabilization in large volume compactifications. Type IIB string theory compactified on Calabi-Yau orientifolds yields $\mathcal{N} = 1$ supergravity as the four dimensional effective theory. Ignoring the gauge sectors, the theory is

specified by the Kähler potential and the superpotential. Including the leading α' corrections [9], the Kähler potential is the form of

$$K = -\log \left[-i \int_M \Omega \wedge \bar{\Omega} \right] - \log [-i(\tau - \bar{\tau})] - 2 \log \left[\frac{\xi}{2} \left(\frac{-i(\tau - \bar{\tau})}{2} \right)^{3/2} + e^{-3\phi_0/2} \mathcal{V} \right], \quad (1)$$

where τ is the axion-dilaton field, Ω is the (3,0)-form of the Calabi-Yau, $\mathcal{V}$ is the classical volume of M in units of $l_s = (2\pi)\sqrt{\alpha'}$, and $\xi = -\zeta(3)\chi(M)/(2(2\pi)^3)$. We require that $\xi > 0$, or $h^{2,1} > h^{1,1}$. Even though the superpotential W [10] does not receive any α' corrections, it acquires a non-perturbative part due to D3-brane instantons [11] or gaugino condensation from wrapped D7-branes [12]:

$$W = \int_M G_3 \wedge \Omega + \sum_i A_i e^{ia_i \rho_i}. \quad (2)$$

Here $G_3 = F_3 - \tau H_3$, with F_3 and H_3 are RR and NS-NS 3-form fluxes, respectively, A_i 's are one-loop determinants, $a_i = 2\pi/M$, with M is a positive integer. Also, $\rho_i \equiv b_i + i\tau_i$ are the complexified Kähler moduli consisting of the four-cycle moduli τ_i 's:

$$\tau_i = \partial_{t^i} \mathcal{V} = \frac{1}{2} D_{ijk} t^j t^k, \quad (3)$$

and the axions b_i 's. The t^i 's measure the areas of two-cycles, D_{ijk} are the triple intersection numbers of the divisor basis [7] and the classical volume is expressed as

$$\mathcal{V} = \frac{1}{6} D_{ijk} t^i t^j t^k. \quad (4)$$

Since the non-perturbative effects in the superpotential are exponentially small compared to the Gukov-Vafa-Witten part, we can follow KKLT by approaching the problem of moduli stabilization in two steps. First, we turn on the background fluxes to stabilize the dilaton and complex structure moduli. Then, we incorporate the non-perturbative effects and stabilize the Kähler moduli while treating the dilaton and complex structure moduli as fixed.

The Kähler potential and the superpotential now become

$$K = K_{cs} - 2 \log \left[\frac{\xi}{2} \left(\frac{-i(\tau - \bar{\tau})}{2} \right)^{3/2} + e^{-3\phi_0/2} \mathcal{V} \right],$$

$$W = W_0 + \sum_i A_i e^{ia_i \rho_i}, \quad (5)$$

where all but the Kähler moduli are taken to be fixed. From this we calculate the potential to be

$$V \equiv V_{\alpha'} + V_{\text{np1}} + V_{\text{np2}},$$

$$V_{\alpha'} = 3\xi e^K \frac{\xi^2 + 7\xi\mathcal{V} + \mathcal{V}^2}{(\mathcal{V} - \xi)(2\mathcal{V} + \xi)^2} |W|^2,$$

$$V_{\text{np1}} = e^K G^{\rho_j \bar{\rho}_k} \left(a_j A_j a_k \bar{A}_k e^{i(a_j \rho_j - a_k \bar{\rho}_k)} \right),$$

$$V_{\text{np2}} = e^K i(a_j A_j e^{ia_j \rho_j} \bar{W} \partial_{\bar{\rho}_k} K - a_k \bar{A}_k e^{-ia_k \bar{\rho}_k} W \partial_{\rho_j} K). \quad (6)$$

Let us consider the large volume limit where all $\tau_i \rightarrow \infty$ except one, which we denote by τ_s . One of the conditions for τ_s is that this limit should be well defined. In this limit, the potential reduces to

$$V = V'_{\alpha'} + V'_{\text{np1}} + V'_{\text{np2}} + V_{\text{supp}},$$

$$V'_{\alpha'} \sim \frac{\xi}{\mathcal{V}^3} e^{K_{cs}} |W_0|^2,$$

$$V'_{\text{np1}} = e^K G^{\rho_s \bar{\rho}_s} a_s^2 |A_s|^2 e^{-2a_s \tau_s}$$

$$\sim \frac{(-k_{ssj} t^j) a_s^2 |A_s|^2 e^{-2a_s \tau_s} e^{K_{cs}}}{\mathcal{V}},$$

$$V'_{\text{np2}} = e^K G^{\rho_s \bar{\rho}_j} i a_s (A_s e^{ia_s \rho_s} \bar{W} \partial_{\bar{\rho}_k} K - \bar{A}_s e^{-ia_s \bar{\rho}_s} W \partial_{\rho_k} \bar{K}). \quad (7)$$

Here, $G_{\rho_s \bar{\rho}_j}$ is the Kähler metric on the size moduli space and V_{supp} denotes the exponentially-suppressed contributions to the potential. Since $V - V_{\text{supp}}$ depends only on b_s , τ_s , and $\mathcal{V}$, we can take advantage of the hierarchy in scales and approach the problem of stabilizing Kähler moduli in two steps, just as in the KKLT construction. Specifically, after fixing the dilaton and complex structure moduli, instead of turning on all the non-perturbative effects, we first only turn on τ_s -dependent effects. Once b_s , τ_s , and $\mathcal{V}$ are stabilized, we treat them as fixed and go back to V_{supp} to stabilize the rest of the Kähler moduli.

The only dependence on the axion b_s is in V'_{np2} . Furthermore, extremizing the potential with respect to b_s will result in making V'_{np2} negative. BBCQ then showed that the potential approaches zero from below for large $\mathcal{V}$ so that there is a large-volume Anti de Sitter minimum. They also found that at this minimum τ_s is stabilized at a small value, which goes along with the earlier assumption of large volume limit.

This solution can be then uplifted to a de Sitter minimum by adding anti-D3-branes [2] or by using the supersymmetric D-terms [13]. The uplift potential is of the

form

$$V_{\text{uplift}} = \frac{\epsilon_{\text{uplift}}}{\mathcal{V}^3}, \quad (8)$$

where $\epsilon_{\text{uplift}} \geq 0$.

We are now ready to try to find the inflaton amongst all the moduli. Consider Calabi-Yau orientifolds with two Kähler moduli ρ_s and ρ_l .

After stabilizing b_s , τ_s , and $\mathcal{V}$, the potential reduces to

$$V = V_0 + V_{\text{supp}}, \quad (9)$$

where V_0 is a constant. Since $\mathcal{V} = \mathcal{V}(\tau_s, \tau_l)$, this means that τ_l has also been stabilized and so $V_{\text{supp}} = V_{\text{supp}}(b_l)$ and

$$V_{\text{supp}} = \frac{e^K G^{\rho_l \bar{\rho}_j} a_l t_j e^{-a_l \tau_l}}{2\mathcal{V}} (A_l \bar{W}_0 e^{ia_l b_l} + \bar{A}_l W_0 e^{-ia_l b_l})$$

$$= \frac{e^K G^{\rho_l \bar{\rho}_j} a_l t_j e^{-a_l \tau_l}}{\mathcal{V}} |A_l W_0| \cos(\theta_W - a_l b_l), \quad (10)$$

where $\bar{A}_l W_0 \equiv |A_l W_0| e^{i\theta_W}$.

The range of V_0 is model-dependent. However, for $\epsilon_{\text{uplift}} = 0$, $V_0 < 0$. Furthermore, for some values $\epsilon_{\text{uplift}} > 0$, $V_0 > 0$, though in this case, ϵ_{uplift} must be sufficiently small such that the minimum is not entirely wiped out. Since the value of V_0 is a continuous function of ϵ_{uplift} , one should be able to tune the parameters to get V_0 to be exactly zero regardless of the geometry. Assuming that this has been done, we then proceed by setting $\theta_W = 0$ and get $V = (\text{constant}) \cos(a_l b_l)$.

As is typical in supergravity theories, the kinetic term for the axion b_l is not canonical. However, since we have fixed the other moduli, we can rescale $b_l \rightarrow \lambda b_l$ with $\lambda = \sqrt{2/G_{\rho_l \bar{\rho}_l}}$, which is constant, to make the kinetic term canonical. The equation of motion in an FRW background is:

$$\ddot{b}_l + 3H\dot{b}_l + \frac{\partial V}{\partial b_l} = 0; \quad 3H^2 = \frac{1}{2}\dot{b}_l^2 + V(b_l), \quad (11)$$

where dots denote cosmic time derivative and in our units, $8\pi G_N = 1$.

The slow-roll parameters are given by [14]

$$\epsilon = \frac{1}{2} \left(\frac{\partial V / \partial b_l}{V} \right)^2 = \frac{a_l^2}{2\lambda^2} \tan^2 \left(\frac{a_l b_l}{\lambda} \right),$$

$$\eta = \frac{\partial^2 V / \partial^2 b_l}{V} = -\frac{a_l^2}{\lambda^2}. \quad (12)$$

Notice that η is constant, which means that once we get a small η , the validity of slow-roll is solely determined by ϵ . Of course, this is not true for $V_0 \neq 0$.

We can rewrite the equations of motion in terms of the number of e -folds N , where $N = \int H dt$:

$$H^2 = \frac{V}{3 - \frac{1}{2}b_l'^2},$$




FIG. 1: $H = \frac{\dot{a}}{a}$, where a is the scale factor, is constant. Thus the scale factor is exponentially increasing.

$$b_l'' + \left(3 - \frac{1}{2}b_l'^2\right) b_l' - \frac{a_l}{\lambda} \left(3 - \frac{1}{2}b_l'^2\right) \tan\left(\frac{a_l b_l}{\lambda}\right) = 0. \quad (13)$$

Using $\eta = -a_l^2/\lambda^2$, we can rewrite ϵ as a function of η

$$\epsilon = \frac{1}{2} |\eta| \tan^2\left(\sqrt{|\eta|} b_l\right). \quad (14)$$

To get inflation, we need to tune η to be small and let the axion b_l roll from near the top of the potential. Now λ depends on the geometry. However, it is constant and since $a = 2\pi/M$, where $M \in \mathbb{Z}^+$ can be a large integer, tuning η is not a serious problem. It is interesting that our ability to get a small η comes mainly from the *discrete* parametrization of the gauge field $SU(M)$ on the wrapped D7-brane.

Furthermore, we have a constraint from the observation of the scalar power spectrum, namely that the spectral index $n_s = 0.951^{+0.015}_{-0.019}$ [15]. For slow-roll inflation, we have the constraint [14]

$$n - 1 = 2\eta - 6\epsilon = -|\eta| \left(2 + 3 \tan^2\left(\sqrt{|\eta|} b_l\right)\right) < 0. \quad (15)$$

This tells us that we need $|\eta| \lesssim 0.03$. For concreteness, $\eta = -0.006$. If we take as initial conditions $b_l(N=0) = 0.1$ and $b_l'(N=0) = 0$, we get 800 e-folds or so of inflation (Fig.1).

The spectral index for the last 60 e-folds before the end of inflation is plotted in Fig.2, and we can see that the observational constraint is compatible with our results.

Since V_{supp} is exponentially suppressed, the scale of inflation is far below the 3×10^{16} GeV threshold required to produce observable tensor contribution to the power spectrum.

We can get sufficient inflation in this model for a large range of initial values of the axion field b_l , although they have to be relatively near the top of the potential. How likely this is to happen requires further analysis.




FIG. 2: Scalar spectral index from axionic inflation for the last 60 e-folds before the end of inflation. The shaded region corresponds to the bounds from WMAP3.

String theory gives us a large number of scalar fields, one or more of which might have been the driving force of inflation in the early universe. However, it is rare to find a single field inflation model from string theory as we have done in this paper. The reason is that in general, at every stage of moduli stabilization, we are dealing with more than one scalar field. At each stage, it is tempting to constrain all fields, except one or two, along the minimum and use the rest, let us call them dynamical field(s), as inflaton(s). For example, when fixing b_s , τ_s , and $\mathcal{V}$, one might want to constrain b_s and $\mathcal{V}$, and only treat τ_s as a dynamical field. However, in general, this is not valid because:

1. Since we have not finished stabilizing all fields in that stage, the constrained fields are still functions of the dynamical fields. For example, before stabilizing τ_s , extremizing the potential with respect to b_s and $\mathcal{V}$ only resulted in expressing them as functions of τ_s , but not as fixed values. Thus, as the dynamical fields evolve, the constrained fields evolve too. Therefore, since all fields are coupled to gravity, in general, we can not neglect the dynamics of the constrained fields in doing the analysis for inflation.
2. We cannot assume *a priori* that as the dynamical fields evolve, the constrained fields will remain constrained. It is more likely that these fields will oscillate around their respective constrained forms. Whether the oscillation can be neglected requires careful analysis of the equations of motion for the constrained fields.

What is different in our situation is that we have used the fact that V_{supp} is exponentially suppressed to approach the problem of moduli stabilization in three stages, where in the last stage we only have to deal with the axion b_l , which we then use as the inflaton.

In order to do a geometry-independent argument, we have tuned V_0 to be zero. As we have discussed above, this is not an unnatural thing to do from the perspective of flux compactifications. Furthermore, $V_0 = 0$ is not a necessary condition for inflation either. We can still produce inflation with $V_0 < 0$ as long as the top of potential is still above zero. Obviously, $V_0 > 0$ will not prevent inflation from happening. However, in order for inflation to stop, V_0 should be sufficiently small such that the bottom of the potential is not above zero.

The theory of reheating is still an open problem in this model. Since we are not working on a specific geometry, we are not going to discuss the problem of reheating in this paper. However, with the recent progress in the study of phenomenological implications in particle physics that emerge from large volume compactifications, esp. concerning mechanism of supersymmetry breaking [16], it would be interesting to work on a specific model that has both inflationary sector and SUSY breaking sector, and address the issue of reheating. We hope to do so in a future publication.

ACKNOWLEDGEMENT

We would like to thank C. P. Burgess for helpful discussions. This work was supported in part by DOE grant DE-FG03-91-ER40682.

* Electronic address: rh4a@andrew.cmu.edu

† Electronic address: jhutasoi@andrew.cmu.edu

- [1] S. B. Giddings, S. Kachru and J. Polchinski, *Hierarchies from fluxes in string compactifications*, Phys. Rev. **D66**, 106006 (2002) [arXiv:hep-th/0105097].
- [2] S. Kachru, R. Kallosh, A. Linde and S. P. Trivedi, *De Sitter Vacua in String Theory* Phys. Rev. D **68**, 046005 (2003) [arXiv:hep-th/0403203].
- [3] M. Grana, *Flux Compactifications in String Theory: A Comprehensive Review*, Phys. Rept. **423**, 91-158 (2006) [arXiv:hep-th/0509003].
- [4] C. P. Burgess, *Inflationary String Theory?*, hep-th/0408037.
- [5] J. J. Blanco-Pillado *et al.*, *Racetrack Inflation*, J. High Energy Phys. **0411**, 063 (2004) [arXiv:hep-th/0406230]; J. J. Blanco-Pillado *et al.* *Inflating in a better Racetrack*, hep-th/0603129.
- [6] V. Balasubramanian, P. Berglund, J.P. Conlon and F. Quevedo, *Systematics of Moduli Stabilization in Calabi-Yau Flux Compactifications*, J. High Energy Phys **0503**, 007 (2005) [arXiv:hep-th/0502058].
- [7] F. Denef, M.R. Douglas and B. Florea, *Building a better Racetrack* J. High Energy Phys. **0406**, 034 (2004) [arXiv:hep-th/0404257].
- [8] K. Freese, J. A. Frieman and A. V. Olinto, *Natural Inflation with Pseudo-Nambu-Goldstone Bosons*, Phys. Rev. Lett. **65**, 3233 (1990); F. C. Adams, J. R. Bond, K. Freese, J. A. Frieman and A. V. Olinto, *Natural Inflation: Particle Physics Models, Power Law Spectra for Large Scale Structure, and Constraints from COBE*, Phys. Rev. D **47**, 426 (1993) [arXiv:hep-ph/9207245].
- [9] K. Becker, M. Becker, M. Haack and J. Louis, *Supersymmetry Breaking and α' -corrections to Flux Induced Potentials*, J. High Energy Phys **0206**, 060 (2002) [arXiv:hep-th/0204254].
- [10] S. Gukov, C. Vafa and E. Witten, *CFT's from Calabi-Yau Fourfolds*, Nucl. Phys. **B584**, 002 (2000). [Erratum-ibid. **B608**, 477 (2001)] [arXiv:hep-th/9906070].
- [11] E. Witten, *Non-Perturbative Superpotentials in String Theory*, Nucl. Phys. **B474**, 343 (1996) [arXiv:hep-th/9604030].
- [12] See for example: C.P. Burgess, J.P. Derendinger, F. Quevedo and M. Quirós, *Gaugino Condensates and Chiral-Linear Duality: An Effective-Lagrangian Analysis*, Phys. Letter B **348**, 428-442 (1995).
- [13] C.P. Burgess, R. Kallosh and F. Quevedo, *de Sitter String Vacua from Supersymmetric D-terms*, J. High Energy Phys. **0310**, 056 (2003) [arXiv:hep-th/0309187].
- [14] A. Riotto, *Inflation and the Theory of Cosmological Perturbations*, hep-ph/0210162.
- [15] Spergel *et al.*, *Wilkinson Microwave Anisotropy Probe (WMAP) Three Year Results: Implications for Cosmology*, <http://lambda.gsfc.nasa.gov/product/map> (2006).
- [16] B. C. Allanach, F. Quevedo and K. Suruliz, *Low-Energy Supersymmetry Breaking from String Flux Compactifications: Benchmark Scenarios*, hep-ph/0512081.